

Example Exam for Physics 222


Remember that these questions are only intended to give you a feel for what my tests are like. They do not cover all of the material you will be expected to know on the real exam.

Useful Integrals:


If a is a real constant greater than zero, then...

$$\int_{-\infty}^{\infty} e^{-a^2 x^2} dx = \frac{\sqrt{\pi}}{a} \qquad \int_{-\infty}^{\infty} x^2 e^{-a^2 x^2} dx = \frac{1}{2} \frac{\sqrt{\pi}}{a^3}$$
$$\int_{-\infty}^{\infty} x e^{-a^2 x^2} dx = 0 \qquad \int_{-\infty}^{\infty} x^3 e^{-a^2 x^2} dx = 0$$

1. An electron in free space has a wave function at time $t = 0$ which is given by $\Psi(x, t = 0) = A e^{-x^2/a^2}$ where a is a constant. The wave function at time zero is plotted below



- (a) Assuming that A is real, what is A ?
- (b) At time $t = 0$ what is $\langle p \rangle$?
- (c) At time $t = 0$ what is $\langle x \rangle$?
- (d) Is this function an eigenfunction of momentum?
2. The dark line below represents the potential $U(x)$. The energies of the $n = 3$ and $n = 6$ energy eigenstates are shown with dashed lines. Make a sketch of what the stationary state wave functions will look like on the dashed lines. Assume that the $n = 1$ state is the ground state.



3. A lithium atom ($Z = 3$) has been stripped of two of its electrons, such that only one electron remains.
- What is the energy of the ground state of the ion?
 - What is the shortest wavelength of light that the ion can emit?
 - What is the lowest final state n_f that can be involved in the emission of a visible photon? (Remember that visible light includes wavelengths between 380 and 760 nm).
4. The dispersion relation for deep water waves (i.e. waves for which $h \gg \lambda$) is given by the relation $\omega = \sqrt{gk}$ where $g = 9.8 \text{ m/s}^2$ is the acceleration due to gravity. Imagine that you are sitting in a row boat in the middle of Utah lake on a calm day. You grab a paddle, dip it into the water, and swish it quickly back and forth 15 times in the space of 2 seconds, generating a water wave packet which travels towards the shore.
- Find an algebraic expression in terms of g and the average wavenumber k for the speed at which the individual ripples within the wave packet travel.
 - Find an algebraic expression in terms of g and the average wavenumber k for the speed at which the packet as a whole travels towards the shore.
 - Find an algebraic expression for k as a function of the average frequency f and the phase speed v_p .
 - Plug the relation found in (c) into the equations found in (a) and (b) and solve the equations to get a numerical answer for the two velocities found in (a) and (b).
5. A particle in an infinite square well of length L has a wave function at time $t = 0$ which is given by

$$\Psi(x, t = 0) = \sqrt{\frac{1}{3L}} \left[\sqrt{2} \sin\left(\frac{3\pi x}{L}\right) + 2e^{i\pi/6} \sin\left(\frac{2\pi x}{L}\right) \right]$$

- What is the wave function at an arbitrary time t ?
 - If you measured the energy of the particle, what is the probability of finding it in the ground state?
 - What is $\langle E \rangle$ at an arbitrary time t ?
- 6.
- What does the term “zero-point” energy mean?
 - Describe the difference between classical and quantum uncertainty.
 - What is the correspondence principle?
 - Explain how the photoelectric effect led to Einstein’s conclusion that light was quantized.
 - What are two of the properties that a wave function must have in order to be “well behaved.”